

Discharge Planning: Optimize Linkage to SUD Care and Supportive Services

References

1. Byrne KA, Roth PJ, Merchant K, et al. Inpatient link to peer recovery coaching: Results from a pilot randomized control trial. *Drug Alcohol Depend.* 2020;215:108234. doi:10.1016/j.drugalcdep.2020.108234.
2. Englander H, Dobbartin K, Lind BK, et al. Inpatient addiction medicine consultation and post-hospital substance use disorder treatment engagement: A propensity-matched analysis. *J Gen Intern Med.* 2019;34(12):2796-2803. doi:10.1007/s11606-019-05251-9.
3. James H, Morgan J, Ti L, Nolan S. Transitions in care between hospital and community settings for individuals with a substance use disorder: A systematic review. *Drug Alcohol Depend.* 2023;243(May 2022):109763. doi:10.1016/j.drugalcdep.2023.109763.
4. Kaiser Permanente Washington Health Research Institute. Structural Vulnerability Assessment Tool. *Systematic Review of Social Risk Screening Tools*. Published June 2020. Accessed March 6, 2025. <https://sdh-tools-review.kpWASHINGTONresearch.org/screening-tools/structural-vulnerability-assessment-tool>.
5. Khalid L, Cunningham CO, Deng Y, et al. Cascade of care for office-based buprenorphine treatment in Bronx community clinics. *J Subst Abuse Treat.* 2022;139(March):108778. doi:10.1016/j.jsat.2022.108778.
6. Liebschutz JM, Crooks D, Herman D, et al. Buprenorphine treatment for hospitalized, opioid-dependent patients. *JAMA Intern Med.* 2014;174(8):1369. doi:10.1001/jamainternmed.2014.2556.
7. Nordeck CD, Welsh C, Schwartz RP, Mitchell SG, O'Grady KE, Gryczynski J. Opioid agonist treatment initiation and linkage for hospitalized patients seen by a substance use disorder consultation service. *Drug Alcohol Depend Reports.* 2022;2(February):100031. doi:10.1016/j.dadr.2022.100031.
8. Perera R, Stephan L, Appa A, et al. Meeting people where they are: Implementing hospital-based substance use harm reduction. *Harm Reduct J.* 2022;19(1):1-7. doi:10.1186/s12954-022-00594-9.
9. Roy PJ, Price R, Choi S, et al. Shorter outpatient wait-times for buprenorphine are associated with linkage to care post-hospital discharge. *Drug Alcohol Depend.* 2021;224(October 2020):108703. doi:10.1016/j.drugalcdep.2021.108703.
10. Wakeman SE, Metlay JP, Chang Y, Herman GE, Rigotti NA. Inpatient addiction consultation for hospitalized patients increases post-discharge abstinence and reduces

addiction severity. *J Gen Intern Med*. 2017;32(8):909-916. doi:10.1007/s11606-017-4077-z.